

# On series of translates of positive functions III

Zoltán Buczolic<sup>\*</sup>, Department of Analysis, ELTE Eötvös Loránd University, Pázmány Péter Sétány 1/c, 1117 Budapest, Hungary

email: [buczo@cs.elte.hu](mailto:buczo@cs.elte.hu)

[www.cs.elte.hu/~buczo](http://www.cs.elte.hu/~buczo)

ORCID Id: 0000-0001-5481-8797

Balázs Maga<sup>†</sup>, Department of Analysis, ELTE Eötvös Loránd University, Pázmány Péter Sétány 1/c, 1117 Budapest, Hungary

email: [magab@cs.elte.hu](mailto:magab@cs.elte.hu)

[www.cs.elte.hu/~magab](http://www.cs.elte.hu/~magab)

and

Gáspár Vertesy<sup>‡</sup>, Department of Analysis, ELTE Eötvös Loránd University, Pázmány Péter Sétány 1/c, 1117 Budapest, Hungary

email: [vertesy.gaspar@gmail.com](mailto:vertesy.gaspar@gmail.com)

January 30, 2018

---

<sup>\*</sup>Research supported by the Hungarian National Research, Development and Innovation Office–NKFIH, Grant 124003.

<sup>†</sup>This author was supported by the ÚNKP-17-2 New National Excellence of the Hungarian Ministry of Human Capacities, and by the Hungarian National Research, Development and Innovation Office–NKFIH, Grant 124003.

<sup>‡</sup>This author was supported by the Hungarian National Research, Development and Innovation Office–NKFIH, Grant 124749.

*Mathematics Subject Classification:* Primary : 28A20, Secondary : 40A05.

*Keywords:* almost everywhere convergence, asymptotically dense, Borel–Cantelli lemma.

*Dedicated to the memory of Jean-Pierre Kahane*

### Abstract

Suppose  $\Lambda$  is a discrete infinite set of nonnegative real numbers. We say that  $\Lambda$  is of type 1 if the series  $s(x) = \sum_{\lambda \in \Lambda} f(x + \lambda)$  satisfies a zero-one law. This means that for any non-negative measurable  $f : \mathbb{R} \rightarrow [0, +\infty)$  either the convergence set  $C(f, \Lambda) = \{x : s(x) < +\infty\} = \mathbb{R}$  modulo sets of Lebesgue zero, or its complement the divergence set  $D(f, \Lambda) = \{x : s(x) = +\infty\} = \mathbb{R}$  modulo sets of measure zero. If  $\Lambda$  is not of type 1 we say that  $\Lambda$  is of type 2.

In this paper we show that there is a universal  $\Lambda$  with gaps monotone decreasingly converging to zero such that for any open subset  $G \subset \mathbb{R}$  one can find a characteristic function  $f_G$  such that  $G \subset D(f_G, \Lambda)$  and  $C(f_G, \Lambda) = \mathbb{R} \setminus G$  modulo sets of measure zero.

We also consider the question whether  $C(f, \Lambda)$  can contain non-degenerate intervals for continuous functions when  $D(f, \Lambda)$  is of positive measure.

The above results answer some questions raised in a paper of Z. Buczolich, J-P. Kahane, and D. Mauldin.

## 1 Introduction

This paper was written for the Kahane memorial volume of Analysis Mathematica. We selected a topic related to Jean-Pierre Kahane's work and decided to answer some questions raised in paper [1] by Z. Buczolich, J-P. Kahane, and D. Mauldin.

This line of research was started in another joint paper with Dan Mauldin [3]. In that paper we considered a problem from 1970, originating from the Diplomarbeit of Heinrich von Weizsäcker [8].

*Suppose  $f : (0, +\infty) \rightarrow \mathbb{R}$  is a measurable function. Is it true that  $\sum_{n=1}^{\infty} f(nx)$  either converges (Lebesgue) almost everywhere or diverges almost everywhere, i.e. is there a zero-one law for  $\sum f(nx)$ ?*

This question also appeared in a paper of J. A. Haight [5].

In [5] it was proved that there exists a set  $H \subset (0, \infty)$  of infinite measure, for which for all  $x, y \in H$ ,  $x \neq y$  the ratio  $x/y$  is not an integer, and furthermore

(†) *for all  $x > 0$   $nx \notin H$  if  $n$  is sufficiently large.*

This implies that if  $f(x) = \chi_H(x)$ , the characteristic function of  $H$  then  $\int_0^{\infty} f(x) dx = \infty$  and  $\sum_{n=1}^{\infty} f(nx) < \infty$  everywhere.

Lekkerkerker in [7] started to study sets with property (†).

In [3] we answered the Haight–Weizsäcker problem.

**Theorem 1.1.** *There exists a measurable function  $f : (0, +\infty) \rightarrow \{0, 1\}$  and two nonempty intervals  $I_F, I_{\infty} \subset [\frac{1}{2}, 1)$  such that for every  $x \in I_{\infty}$  we have  $\sum_{n=1}^{\infty} f(nx) =$*

$+\infty$  and for almost every  $x \in I_F$  we have  $\sum_{n=1}^{\infty} f(nx) < +\infty$ . The function  $f$  is the characteristic function of an open set  $E$ .

Jean-Pierre Kahane was interested in this problem and soon after our paper had become available we started to receive faxes and emails from him. This cooperation lead to papers [1] and [2].

We considered a more general, additive version of the Haight–Weizsäcker problem. Since  $\sum_{n=1}^{\infty} f(nx) = \sum_{n=1}^{\infty} f(e^{\log x + \log n})$ , that is using the function  $h = f \circ \exp$  defined on  $\mathbb{R}$  and  $\Lambda = \{\log n : n = 1, 2, \dots\}$  we were interested in almost everywhere convergence questions of the series  $\sum_{\lambda \in \Lambda} h(x + \lambda)$ .

Taking more general sets than  $\Lambda = \{\log n : n = 1, 2, \dots\}$  was also motivated by a paper, [6] of Haight. He proved, using the original multiplicative notation of our problem that if  $\Lambda \subset [0, +\infty)$  is an arbitrary countable set such that its only accumulation point is  $+\infty$  then there exists a measurable set  $E \subset (0, +\infty)$  of infinite measure such that for all  $x, y \in E$ ,  $x \neq y$ ,  $x/y \notin \Lambda$ , and for a fixed  $x$  there exist only finitely many  $\lambda \in \Lambda$  for which  $\lambda x \in E$ . This implies that choosing  $f = \chi_E$  we have  $\sum_{\lambda \in \Lambda} f(\lambda x) < \infty$ , but  $\int_{\mathbb{R}^+} f(x) dx = \infty$ .

Next we recall from [1] the definition of type 1 and type 2 sets. Given  $\Lambda$  an unbounded, infinite discrete set of nonnegative numbers, and a measurable  $f : \mathbb{R} \rightarrow [0, +\infty)$ , we consider the sum

$$s(x) = \sum_{\lambda \in \Lambda} f(x + \lambda),$$

and the complementary subsets of  $\mathbb{R}$ :

$$C = C(f, \Lambda) = \{x : s(x) < \infty\}, \quad D = D(f, \Lambda) = \{x : s(x) = \infty\}.$$

**Definition 1.2.** The set  $\Lambda$  is of type 1 if, for every  $f$ , either  $C(f, \Lambda) = \mathbb{R}$  a.e. or  $C(f, \Lambda) = \emptyset$  a.e. (or equivalently  $D(f, \Lambda) = \emptyset$  a.e. or  $D(f, \Lambda) = \mathbb{R}$  a.e.). Otherwise,  $\Lambda$  has type 2.

That is for type 1 sets we have a "zero-one" law for the almost everywhere convergence properties of the series  $\sum_{\lambda \in \Lambda} f(x + \lambda)$ , while for type 2 sets the situation is more complicated.

**Definition 1.3.** The unbounded, infinite discrete set  $\Lambda = \{\lambda_1, \lambda_2, \dots\}$ ,  $\lambda_1 < \lambda_2 < \dots$  is asymptotically dense if  $d_n = \lambda_n - \lambda_{n-1} \rightarrow 0$ , or equivalently:

$$\forall a > 0, \quad \lim_{x \rightarrow \infty} \#(\Lambda \cap [x, x + a]) = \infty.$$

If  $d_n$  tends to zero monotone decreasingly, we speak about decreasing gap asymptotically dense sets.

If  $\Lambda$  is not asymptotically dense we say that it is asymptotically lacunary.

We denote the non-negative continuous functions on  $\mathbb{R}$  by  $C^+(\mathbb{R})$ , and if, in addition these functions tend to zero in  $+\infty$  they belong to  $C_0^+(\mathbb{R})$ .

In [1] we gave some necessary and some sufficient conditions for a set  $\Lambda$  being of type 2. A complete characterization of type 2 sets is still unknown. We recall here from [1] the theorem concerning the Haight–Weizsäcker problem. This contains the additive version of the result of Theorem 1.1 with some additional information.

**Theorem 1.4.** *The set  $\Lambda = \{\log n : n = 1, 2, \dots\}$  has type 2. Moreover, for some  $f \in C_0^+(\mathbb{R})$ ,  $C(f, \Lambda)$  has full measure on the half-line  $(0, \infty)$  and  $D(f, \Lambda)$  contains the half-line  $(-\infty, 0)$ . If for each  $c$ ,  $\int_c^{+\infty} e^y g(y) dy < +\infty$ , then  $C(g, \Lambda) = \mathbb{R}$  a.e. If  $g \in C_0^+(\mathbb{R})$  and  $C(g, \Lambda)$  is not of the first (Baire) category, then  $C(g, \Lambda) = \mathbb{R}$  a.e. Finally, there is some  $g \in C_0^+(\mathbb{R})$  such that  $C(g, \Lambda) = \mathbb{R}$  a.e. and  $\int_0^{+\infty} e^y g(y) dy = +\infty$ .*

As  $\Lambda$  used in the above theorem is a decreasing gap asymptotically dense set and quite often it is much easier to construct examples with lacunary  $\Lambda$ s, in our paper we try to give examples with a decreasing gap asymptotically dense  $\Lambda$ .

One might believe that for type 2  $\Lambda$ s  $C(f, \Lambda)$ , or  $D(f, \Lambda)$  are always half-lines if they differ from  $\mathbb{R}$ . Indeed in [1] we obtained results in this direction. A number  $t > 0$  is called a translator of  $\Lambda$  if  $(\Lambda + t) \setminus \Lambda$  is finite. Condition (\*) is said to be satisfied if  $T(\Lambda)$ , the countable additive semigroup of translators of  $\Lambda$ , is dense in  $\mathbb{R}^+$ . We showed that condition (\*) implies that  $C(f, \Lambda)$  is either  $\emptyset$ ,  $\mathbb{R}$ , or a right half-line modulo sets of measure zero.

In [4] we showed that this is not always the case. For a given  $\alpha \in (0, 1)$  and a sequence of natural numbers  $n_1 < n_2 < \dots$  we put  $\Lambda^{\alpha^k} := \cup_{k=1}^{\infty} \Lambda_k^{\alpha^k}$ ,  $\Lambda_k^{\alpha^k} = \alpha^k \mathbb{Z} \cap [n_k, n_{k+1})$ .

If  $\alpha = \frac{1}{q}$  for some  $q \in \{2, 3, \dots\}$ , then a slight modification of the proof of Theorem 1 of [1] shows that  $\Lambda^{(\frac{1}{q})^k}$  is of type 1 and condition (\*) is satisfied.

If  $\alpha \notin \mathbb{Q}$ , then one can apply Theorem 5 of [1] to show that  $\Lambda^{\alpha^k}$  is of type 2.

The difficult case is when  $\alpha = \frac{p}{q}$  with  $(p, q) = 1$ ,  $p, q > 1$ ,  $p < q$ . In this case we showed that  $\Lambda^{(\frac{p}{q})^k}$  is of type 2. In the cases  $\Lambda^{(\frac{p}{q})^k}$ , ( $p > 1$ ) condition (\*) is not satisfied and we also showed in [4] that there exists a characteristic function  $f$  such that  $C(f, \Lambda)$  does not equal  $\emptyset$ ,  $\mathbb{R}$ , or a right half-line modulo sets of measure zero. This structure of  $C(f, \Lambda)$  had not been seen before our paper [4].

From the point of view of our current paper the following question (QUESTION 2 in [1]) is the most relevant:

**Question 1.5.** Given open sets  $G_1$  and  $G_2$  when is it possible to find  $\Lambda$  and  $f$  such that  $C(f, \Lambda)$  contains  $G_1$  and  $D(f, \Lambda)$  contains  $G_2$ ?

It was remarked in [1] that if the counting function of  $\Lambda$ ,  $n(x) = \#\{\Lambda \cap [0, x]\}$

satisfies a condition of the type

$$\forall \ell < 0 \forall a \in \mathbb{R} \quad \limsup_{x \rightarrow \infty} \frac{n(x + \ell + a) - n(x + a)}{n(x + \ell) - n(x)} < +\infty$$

(as is the case for  $\Lambda = \{\log n\}$ ) then either  $C(f, \Lambda)$  has full measure on  $\mathbb{R}$  or  $C(f, \Lambda)$  does not contain any interval.

It was also mentioned in [1] that if  $\Lambda$  is asymptotically lacunary then it is possible to construct  $f \in C_0^+(\mathbb{R})$  such that both  $C(f, \Lambda)$  and  $D(f, \Lambda)$  have interior points.

In this paper we give an almost complete answer to Question 1.5. In Section 2 we prove Theorem 2.1. This theorem states that there is a universal decreasing gap asymptotically dense  $\Lambda$  such that for any open subset  $G \subset \mathbb{R}$  one can find a characteristic function  $f_G$  such that  $G \subset D(f_G, \Lambda)$  and  $C(f_G, \Lambda) = \mathbb{R} \setminus G$  modulo sets of measure zero. We also show that one can also select a  $g_G \in C_0^+(\mathbb{R})$  with similar properties.

In Section 3 we consider the question of subintervals in  $C(f, \Lambda)$  when  $f \in C_0^+(\mathbb{R})$ . In Theorem 3.1 we prove that there exists a universal asymptotically dense infinite discrete set  $\Lambda$  such that for any open set  $G \subset \mathbb{R}$  one can select an  $f_G \in C_0^+(\mathbb{R})$  such that  $D(f_G, \Lambda) = G$ . In this case there is no exceptional set of measure zero,  $D(f_G, \Lambda)$  equals  $G$  exactly. On the other hand,  $\Lambda$  is not of decreasing gap. As Theorem 3.4 shows it is impossible to find such a universal  $\Lambda$  with decreasing gaps. In Theorem 3.4 we prove that if  $\Lambda$  is a decreasing gap asymptotically dense set,  $f \in C^+(\mathbb{R})$  and  $x$  is an interior point of  $C(f, \Lambda)$  then  $[x, +\infty) \cap D(f, \Lambda)$  is of zero Lebesgue measure.

The example provided in Theorem 3.3 demonstrates that there is a decreasing gap asymptotically dense  $\Lambda$  and an  $f \in C_0^+(\mathbb{R})$  such that  $D(f, \Lambda)$  and  $C(f, \Lambda)$  both contain interior points. Of course, as Theorem 3.4 shows the interior points of  $D(f, \Lambda)$  are to the left of those of  $C(f, \Lambda)$ .

## 2 A universal decreasing gap asymptotically dense $\Lambda$ set

Let  $\mu$  denote the one-dimensional Lebesgue measure.

We denote by  $\mathbb{N} := \{n \in \mathbb{Z} : n \geq 1\}$  the set of natural numbers. For every  $A, B \subset \mathbb{R}$  we put  $A + B := \{a + b : a \in A \text{ and } b \in B\}$  and  $A - B := \{a - b : a \in A \text{ and } b \in B\}$ .

The integer, and fractional parts of  $x \in \mathbb{R}$  are denoted by  $[x]$  and  $\{x\}$ , respectively.

**Theorem 2.1.** *There is a strictly monotone increasing unbounded sequence  $(\lambda_0, \lambda_1, \dots) = \Lambda$  in  $\mathbb{R}$  such that  $\lambda_n - \lambda_{n-1}$  tends to 0 monotone decreasingly, that is  $\Lambda$  is a decreasing gap asymptotically dense set, such that for every open set  $G \subset \mathbb{R}$  there is a function  $f_G : \mathbb{R} \rightarrow [0, +\infty)$  for which*

$$\mu \left( \left\{ x \notin G : \sum_{n=0}^{\infty} f_G(x + \lambda_n) = \infty \right\} \right) = 0, \text{ and} \quad (1)$$

$$\sum_{n=0}^{\infty} f_G(x + \lambda_n) = \infty \text{ for every } x \in G, \quad (2)$$

moreover  $f_G = \chi_{U_G}$  for a closed set  $U_G \subset \mathbb{R}$ . By (1) and (2) we have  $D(f_G, \Lambda) \supset G$ , and  $C(f_G, \Lambda) = \mathbb{R} \setminus G$  modulo sets of measure zero.

One can also select a  $g_G \in C_0^+(\mathbb{R})$  satisfying (1) and (2) instead of  $f_G$ .

**Remark 2.2.** Observe that in the above theorem we construct a universal  $\Lambda$  and for this set, depending on our choice of  $G$  we can select a suitable  $f_G$  such that  $D(f_G, \Lambda) = G$  modulo sets of measure zero.

*Proof.* Let

$$\mathcal{I} := \{(j, k) : j \in \mathbb{N} \text{ and } k \in \mathbb{Z} \cap [0, 2j \cdot 2^j)\}$$

with the following lexicographical ordering: if  $(j, k), (\tilde{j}, \tilde{k}) \in \mathcal{I}$  then

$$(j, k) <_{\mathcal{I}} (\tilde{j}, \tilde{k}) \Leftrightarrow (j < \tilde{j} \text{ or } (j = \tilde{j} \text{ and } k < \tilde{k})).$$

Given  $(j, k) \in \mathcal{I}$  we define its immediate successor  $(\hat{j}, \hat{k})$  the following way: let  $\hat{j} := j$  and  $\hat{k} := k + 1$  if  $k < 2j \cdot 2^j - 1$ , and let  $\hat{j} := j + 1$  and  $\hat{k} := 0$  if  $k = 2j \cdot 2^j - 1$ . It is clear that starting with  $(1, 0)$  by repeated application of taking the immediate successor we can enumerate  $\mathcal{I}$  and hence we will be able to do induction on  $\mathcal{I}$ . We will also introduce the operation of taking the predecessor of  $(j, k) \neq (1, 0)$  which will be denoted by  $(\check{j}, \check{k})$  and which is defined by the property  $(\hat{\check{j}}, \hat{\check{k}}) = (j, k)$ .


For every  $(j, k) \in \mathcal{I}$  let

$$I_{j,k} := [j - (k + 1)2^{-j}, j - k2^{-j}] = [a_{I_{j,k}}, b_{I_{j,k}}].$$

In (6) a set  $U_{j,k}$  will be defined such that with a properly selected  $\Lambda$  we have

$$I_{j,k} \subset U_{j,k} - \Lambda = \{x \in \mathbb{R} : \exists n \in \mathbb{N} \cup \{0\} \text{ such that } x + \lambda_n \in U_{j,k}\} \text{ and} \quad (3)$$

$$\begin{aligned} \mu(\{x \in [-j, j] : \exists \text{ infinitely many } (j^*, k^*) \in \mathcal{I} \\ \text{for which } x \in (U_{j^*, k^*} - \Lambda) \setminus I_{j^*, k^*}\}) = 0. \end{aligned} \quad (4)$$



**Figure 1:** Definition of  $I_{j,k}$  and  $U_{j,k}$

Let  $G$  be an arbitrary open subset of  $\mathbb{R}$  and let

$$U_G := \bigcup \{U_{j^*, k^*} : (j^*, k^*) \in \mathcal{I} \text{ and } I_{j^*, k^*} \subset G\}.$$

Put

$$f_G(x) := \begin{cases} 1 & \text{if } x \in U_G \\ 0 & \text{else.} \end{cases} \quad (5)$$

We will prove that  $\Lambda$  and  $f_G$  satisfy the conditions of the theorem.

Now we define the sets  $U_{j,k}$ . Before doing this we recall and introduce some notation. For every  $(j, k) \in \mathcal{I}$  let

- $a_{I_{j,k}} := j - (k + 1) \cdot 2^{-j}$  (that is  $a_{I_{j,k}}$  is the left endpoint of  $I_{j,k}$ ),
- $b_{I_{j,k}} := j - k \cdot 2^{-j}$  (that is  $b_{I_{j,k}}$  is the right endpoint of  $I_{j,k}$ ),
- $E_{j,k} := 2^{-2j \cdot 2^j - k}$ ,
- $a_{j,k} := 2^{2j \cdot 2^j + k}$ ,
- $b_{j,k} := a_{j,k} + E_{j,k}$ .

See Figure 1. This and the other figure in this paper are to illustrate concepts and they are not drawn to illustrate a certain step, for example with a fixed  $j$  of our construction.

Let

$$U_{j,k} := \bigcup_{i=0}^{E_{j,k}^{-1}-1} [a_{j,k} + iE_{j,k}^2, a_{j,k} + iE_{j,k}^2 + E_{j,k}^3] \subset [a_{j,k}, b_{j,k}]. \quad (6)$$

Next we prove a useful lemma:

**Lemma 2.3.** For every  $(j, k) \in \mathcal{I}$  we have

$$a_{j,k} \leq \frac{a_{\hat{j},\hat{k}}}{2} \text{ and } E_{j,k} \geq 2E_{\hat{j},\hat{k}}, \quad (7)$$

moreover,

$$E_{j,k}/2 \text{ is an integer multiple of } E_{\hat{j},\hat{k}}. \quad (8)$$

*Proof.* It is enough to prove (7) for  $a_{j,k}$  as  $E_{j,k} = a_{j,k}^{-1}$ .

First suppose that  $k < 2j \cdot 2^j - 1$ , then  $\hat{j} = j$ ,  $\hat{k} = k + 1$  and

$$a_{j,k} = 2^{2j \cdot 2^j + k} = \frac{2^{2j \cdot 2^j + (k+1)}}{2} = \frac{a_{\hat{j},\hat{k}}}{2}. \quad (9)$$

If  $k = 2j \cdot 2^j - 1$  then  $\hat{j} = j + 1$ ,  $\hat{k} = 0$  and

$$a_{j,k} = 2^{2j \cdot 2^j + k} = 2^{2j \cdot 2^j + 2j \cdot 2^j - 1} = 2^{4j \cdot 2^j - 1} = \frac{2^{2^{(j+1)} \cdot 2^{(j+1)}}}{2^{2 \cdot 2^{j+1} + 1}} = \frac{a_{\hat{j},\hat{k}}}{2^{2 \cdot 2^{j+1} + 1}}. \quad (10)$$

Using  $E_{j,k} = a_{j,k}^{-1}$  from (9) and (10) it follows that (8) holds.  $\square$

Next we turn to the definition of  $\Lambda$ .

During the definition of  $\Lambda$  we will use the notation  $d_n := \lambda_n - \lambda_{n-1}$ , in fact, often we will define  $d_n$  and that will provide the value of  $\lambda_n$  given the already defined  $\lambda_{n-1}$ . Let  $\lambda_0 := a_{1,0} - b_{I_{1,0}}$  and  $n_{0,1,0} = 0$ .

Suppose that for a  $(j, k) \in \mathcal{I}$  we have already defined  $n_{0,j,k}$  and  $\lambda_n$  for  $n \leq n_{0,j,k}$ ,  $\lambda_{n_{0,j,k}} = a_{j,k} - b_{I_{j,k}}$  and  $d_{n_{0,j,k}}/E_{j,k}^2$  is a positive integer (or  $n_{0,j,k} = 0$ ). Now we need to do our next step to define these objects for  $(\hat{j}, \hat{k})$ .

**Step  $(\hat{j}, \hat{k})$ .** Let  $n_{1,j,k} := n_{0,j,k} + 2^{-j}E_{j,k}^{-2} + 2E_{j,k}^{-1}$ . For every integer  $n \in [n_{0,j,k} + 1, n_{1,j,k}]$  let  $d_n := E_{j,k}^2 - E_{j,k}^3$ . Thus we have

$$\begin{aligned} \lambda_{n_{1,j,k}} &= \lambda_{n_{0,j,k}} + (2^{-j}E_{j,k}^{-2} + 2E_{j,k}^{-1})(E_{j,k}^2 - E_{j,k}^3) \\ &= a_{j,k} - b_{I_{j,k}} + 2^{-j} - 2^{-j}E_{j,k} + 2E_{j,k} - 2E_{j,k}^2 \\ &= a_{j,k} - a_{I_{j,k}} + 2E_{j,k} - 2^{-j}E_{j,k} - 2E_{j,k}^2 \\ &= b_{j,k} - a_{I_{j,k}} + E_{j,k} - 2^{-j}E_{j,k} - 2E_{j,k}^2 \geq b_{j,k} - a_{I_{j,k}} \end{aligned} \quad (11)$$

and (from the second row of (11))

$$\lambda_{n_{1,j,k}} = a_{j,k} - b_{I_{j,k}} + 2^{-j} - 2^{-j}E_{j,k} + 2E_{j,k} - 2E_{j,k}^2 < a_{j,k} - b_{I_{j,k}} + 1. \quad (12)$$

Since  $a_{j,k} - a_{I_{j,k}} = 2^{2j \cdot 2^j + k} - (j - k \cdot 2^{-j})$  and  $2^{-j}E_{j,k}$  are both integer multiples of  $E_{j,k}^2 = (2^{-2j \cdot 2^j - k})^2$  from the third row of (11) we obtain that

$$\lambda_{n_{1,j,k}} \text{ is an integer multiple of } E_{j,k}^2. \quad (13)$$



By Lemma 2.3 and (12) we have

$$a_{\hat{j},\hat{k}} - b_{I_{\hat{j},\hat{k}}} \geq 2a_{j,k} - (j+1) \geq a_{j,k} + j + 1 > a_{j,k} - b_{I_{j,k}} + 1 > \lambda_{n_{1,j,k}}.$$

We set

$$n_{0,\hat{j},\hat{k}} = n_{1,j,k} + \frac{a_{\hat{j},\hat{k}} - b_{I_{\hat{j},\hat{k}}} - \lambda_{n_{1,j,k}}}{2^{-1}E_{j,k}^2} \quad (14)$$

and

$$d_n = E_{j,k}^2/2 \text{ for every integer } n \in (n_{1,j,k}, n_{0,\hat{j},\hat{k}}]. \quad (15)$$

We obtain by (14)

$$\lambda_{n_{0,\hat{j},\hat{k}}} = \lambda_{n_{1,j,k}} + \frac{(n_{0,\hat{j},\hat{k}} - n_{1,j,k})E_{j,k}^2}{2} = \lambda_{n_{1,j,k}} + a_{\hat{j},\hat{k}} - b_{I_{\hat{j},\hat{k}}} - \lambda_{n_{1,j,k}} = a_{\hat{j},\hat{k}} - b_{I_{\hat{j},\hat{k}}},$$

and by (8),  $d_{n_{0,\hat{j},\hat{k}}} = E_{j,k}^2/2$  is an integer multiple of  $E_{\hat{j},\hat{k}}^2$ , hence (13) implies that

$$\lambda_n \text{ is an integer multiple of } E_{\hat{j},\hat{k}}^2 \text{ for } n \in (n_{1,j,k}, n_{0,\hat{j},\hat{k}}]. \quad (16)$$

Thus we can proceed to the next step. By repeating this procedure we can carry out the above steps for all  $(j, k) \in \mathcal{I}$  and hence we can define  $\Lambda$ .

Now we prove (3). We fix  $(j, k)$  and choose an arbitrary point  $x$  from  $I_{j,k}$ . Let  $n_x$  denote the smallest integer for which

$$x + \lambda_{n_x} > a_{j,k}. \quad (17)$$

Put  $n'_x := n_x + \left\lfloor \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\rfloor$ .

We have  $x \in I_{j,k} \subset [-j, j]$ . From  $x + \lambda_{n_{0,j,k}} = x + a_{j,k} - b_{I_{j,k}}$  it follows that

$$x + \lambda_{n_{0,j,k}} - a_{j,k} = x - b_{I_{j,k}} \leq 0. \quad (18)$$

Therefore,  $n_x > n_{0,j,k}$  and hence

$$d_n \leq d_{n_{0,j,k}+1} = E_{j,k}^2 - E_{j,k}^3 \text{ for every } n \in [n_x, \infty). \quad (19)$$

By minimality of  $n_x$  we have

$$x + \lambda_{n_x} - a_{j,k} \leq d_{n_x} \leq E_{j,k}^2 - E_{j,k}^3. \quad (20)$$

Next we will show that  $x + \lambda_{n'_x} \in U_{j,k}$ . Using (19)

$$0 \leq \left\lfloor \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\rfloor \leq \frac{d_{n_x}}{E_{j,k}^3} \leq \frac{E_{j,k}^2 - E_{j,k}^3}{E_{j,k}^3} = E_{j,k}^{-1} - 1. \quad (21)$$

We also infer

$$\begin{aligned}
x + \lambda_{n'_x} &= x + \lambda_{n_x} + \sum_{n \in (n_x, n'_x]} d_n \leq x + \lambda_{n_x} + \left\lfloor \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\rfloor (E_{j,k}^2 - E_{j,k}^3) \\
&= a_{j,k} + (x + \lambda_{n_x} - a_{j,k}) + \left\lfloor \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\rfloor (E_{j,k}^2 - E_{j,k}^3) \\
&= a_{j,k} + \left\lfloor \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\rfloor E_{j,k}^2 + E_{j,k}^3 \left\{ \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\} \\
&\text{using (21)} \\
&\leq a_{j,k} + (E_{j,k}^{-1} - 1)E_{j,k}^2 + E_{j,k}^3 \leq a_{j,k} + E_{j,k} = b_{j,k}.
\end{aligned} \tag{22}$$

From (11) and (22) we obtain

$$\lambda_{n'_x} \leq b_{j,k} - x \leq b_{j,k} - a_{I_{j,k}} \leq \lambda_{n_{1,j,k}},$$

hence  $n_x, n'_x \leq n_{1,j,k}$ , which means that  $d_n = E_{j,k}^2 - E_{j,k}^3$  for every  $n \in (n_x, n'_x]$ . This implies that the first inequality in (22) is, in fact an equality, that is

$$x + \lambda_{n'_x} = a_{j,k} + \left\lfloor \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\rfloor E_{j,k}^2 + E_{j,k}^3 \left\{ \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\}. \tag{23}$$

Using (21) and (23) we can see that there exists an integer  $i = \left\lfloor \frac{x + \lambda_{n_x} - a_{j,k}}{E_{j,k}^3} \right\rfloor \in [0, E_{j,k}^{-1} - 1]$  such that

$$a_{j,k} + iE_{j,k}^2 \leq x + \lambda_{n'_x} \leq a_{j,k} + iE_{j,k}^2 + E_{j,k}^3$$

that is  $x + \lambda_{n'_x} \in U_{j,k}$ , which implies (3).

We continue with the proof of (4). Suppose  $(\check{j}, \check{k}), (j, k), (\hat{j}, \hat{k}) \in \mathcal{I}$ . Then they are strictly monotone increasing in this order and are adjacent in the lexicographical ordering of  $\mathcal{I}$ . We have by Lemma 2.3 and the third row of (11)

$$\begin{aligned}
j + \lambda_{n_{1,\check{j},\check{k}}} &= j + a_{\check{j},\check{k}} - a_{I_{\check{j},\check{k}}} + 2E_{\check{j},\check{k}} - 2^{-\check{j}}E_{\check{j},\check{k}} - 2E_{\check{j},\check{k}}^2 \\
&< a_{\check{j},\check{k}} + 2j + 1 \leq 2a_{\check{j},\check{k}} \leq a_{j,k},
\end{aligned} \tag{24}$$

that is  $U_{j,k} - \lambda_{n_{1,\check{j},\check{k}}}$  is to the right of  $j$ . By (16),  $\lambda_n/E_{j,k}^2$  is an integer for every  $n \in (n_{1,\check{j},\check{k}}, n_{0,j,k}]$ . Therefore, (24) implies that

$$\begin{aligned}
B_{j,k} &:= [b_{j,k} - \lambda_{n_{0,j,k}}, j] \cap (U_{j,k} - \Lambda) \\
&= [b_{j,k} - \lambda_{n_{0,j,k}}, j] \cap (U_{j,k} - \{\lambda_n : n \in (n_{1,\check{j},\check{k}}, n_{0,j,k}]\}) \\
&\subset [b_{j,k} - \lambda_{n_{0,j,k}}, j] \cap \bigcup_{i \in \mathbb{Z}} [iE_{j,k}^2, iE_{j,k}^2 + E_{j,k}^3].
\end{aligned} \tag{25}$$

Similarly, by using (7)

$$\begin{aligned} -j + \lambda_{n_{0,j,\hat{k}}} &= -j + a_{j,\hat{k}} - b_{I_{j,\hat{k}}} > a_{j,\hat{k}} - (2j + 1) \\ &\geq 2a_{j,k} - (2j + 1) \geq a_{j,k} + E_{j,k} = b_{j,k}, \end{aligned} \quad (26)$$

that is  $U_{j,k} - \lambda_{n_{0,j,\hat{k}}}$  is to the left of  $-j$ . Since by (13) and (15)  $\lambda_n / (E_{j,k}^2/2)$  is an integer for every  $n \in [n_{1,j,k}, n_{0,j,\hat{k}}]$ , (26) implies that

$$\begin{aligned} A_{j,k} &:= [-j, a_{j,k} - \lambda_{n_{1,j,k}}] \cap (U_{j,k} - \Lambda) \\ &= [-j, a_{j,k} - \lambda_{n_{1,j,k}}] \cap \left( U_{j,k} - \{ \lambda_n : n \in [n_{1,j,k}, n_{0,j,\hat{k}}] \} \right) \\ &\subset [-j, a_{j,k} - \lambda_{n_{1,j,k}}] \cap \bigcup_{i \in \mathbb{Z}} [iE_{j,k}^2/2, iE_{j,k}^2/2 + E_{j,k}^3]. \end{aligned} \quad (27)$$

We want to estimate the following expression from above:

$$\begin{aligned} &\mu([-j, j] \cap (U_{j,k} - \Lambda) \setminus I_{j,k}) \\ &\leq \mu(A_{j,k} \cup [a_{j,k} - \lambda_{n_{1,j,k}}, a_{I_{j,k}}] \cup [b_{I_{j,k}}, b_{j,k} - \lambda_{n_{0,j,k}}] \cup B_{j,k}). \end{aligned} \quad (28)$$

By (25) and (27) we have

$$\begin{aligned} &\mu(A_{j,k} \cup B_{j,k}) \\ &\leq \mu\left([-j, j] \cap \left(\bigcup_{i \in \mathbb{Z}} [iE_{j,k}^2/2, iE_{j,k}^2/2 + E_{j,k}^3]\right)\right) \\ &= E_{j,k}^3 \frac{2j}{E_{j,k}^2/2} = 4j \cdot E_{j,k}, \end{aligned} \quad (29)$$

and using the third row of (11)

$$\begin{aligned} \mu([a_{j,k} - \lambda_{n_{1,j,k}}, a_{I_{j,k}}]) &= a_{I_{j,k}} - (a_{j,k} - (a_{j,k} - a_{I_{j,k}} + 2E_{j,k} - 2^{-j}E_{j,k} - 2E_{j,k}^2)) \\ &= 2E_{j,k} - 2^{-j}E_{j,k} - 2E_{j,k}^2 \leq 2E_{j,k}. \end{aligned} \quad (30)$$

Moreover,

$$\mu[b_{I_{j,k}}, b_{j,k} - \lambda_{n_{0,j,k}}] = b_{j,k} - (a_{j,k} - b_{I_{j,k}}) - b_{I_{j,k}} = b_{j,k} - a_{j,k} = E_{j,k}. \quad (31)$$

Writing (29), (30) and (31) into (28) yields

$$\mu([-j, j] \cap (U_{j,k} - \Lambda) \setminus I_{j,k}) \leq (4j + 3) \cdot E_{j,k}. \quad (32)$$

Thus

$$\sum_{(j^*, k^*) \in \mathcal{I}} \mu([-j, j] \cap (U_{j^*, k^*} - \Lambda) \setminus I_{j^*, k^*}) \quad (33)$$

$$\begin{aligned}
&\leq \sum_{\substack{(j^*, k^*) \in \mathcal{I} \\ j^* < j}} \mu([-j, j] \cap (U_{j^*, k^*} - \Lambda) \setminus I_{j^*, k^*}) \\
&+ \sum_{(j^*, k^*) \in \mathcal{I}} \mu([-j^*, j^*] \cap (U_{j^*, k^*} - \Lambda) \setminus I_{j^*, k^*}) \\
&\leq \sum_{\substack{(j^*, k^*) \in \mathcal{I} \\ j^* < j}} 2j + \sum_{(j^*, k^*) \in \mathcal{I}} (4j^* + 3) \cdot E_{j^*, k^*} \\
&\leq 2j \cdot 2j(2^{j-1} + \dots + 1) + \sum_{j^*=1}^{\infty} \sum_{k^*=0}^{2j^* \cdot 2^{j^*} - 1} (4j^* + 3) E_{j^*, k^*} \\
&\leq 4j^2 \cdot 2^j + \sum_{j^*=1}^{\infty} 2j^* \cdot 2^{j^*} (4j^* + 3) 2^{-2j^* \cdot 2^{j^*}} \\
&\leq 4j^2 \cdot 2^j + \sum_{j^*=1}^{\infty} (8(j^*)^2 + 6j^*) 2^{-2j^* \cdot 2^{j^*} + j^*} < \infty,
\end{aligned}$$

which by the Borel–Cantelli lemma implies (4).

Let  $G$  be a fixed open subset of  $\mathbb{R}$ . If  $x \in G$ , then  $\{(j, k) \in \mathcal{I} : x \in I_{j, k} \subset G\}$  is an infinite set, hence according to (3) and (5)

$$\sum_{n=0}^{\infty} f_G(x + \lambda_n) = \infty.$$

If  $x \in \mathbb{R} \setminus G$  and  $\sum_{n=0}^{\infty} f_G(x + \lambda_n) = \infty$ , then  $\{n \in \mathbb{N} : x + \lambda_n \in U_G\}$  is an infinite set, which implies that  $\{(j^*, k^*) \in \mathcal{I} : I_{j^*, k^*} \subset G \text{ and } x \in (U_{j^*, k^*} - \Lambda)\}$  is also infinite, thus (4) implies (1).

Next we see how one can modify  $f_G$  to obtain a  $g_G \in C_0^+(\mathbb{R})$  still satisfying (1) and (2). In [1] there is Proposition 1, which says that one can modify  $f_G$  to obtain a  $g_G \in C_0^+(\mathbb{R})$  such that  $C(f_G, \lambda) = C(g_G, \lambda)$  a.e. and  $D(f_G, \lambda) = D(g_G, \lambda)$  a.e. Since we want to preserve (2) we cannot change  $D(f_G, \lambda)$  by an arbitrary set of measure zero. Hence in the next construction a little extra care is needed.

$$\text{Put } \Lambda_N = \{\lambda \in \Lambda : \lambda \leq 10N\} \text{ and } L_N = \#\Lambda_N. \quad (34)$$

Observe that  $U_G \cap (-\infty, 0] = \emptyset$ ,  $U_G$  does not contain a half-line, and  $U_G \cap [0, N]$  is the union of finitely many disjoint closed intervals for any  $N \in \mathbb{N}$ .

Choose an open  $\tilde{U}_G \supset U_G$  such that it does not contain a half-line, and

$$\mu((\tilde{U}_G \setminus U_G) \cap [N-1, N]) < \frac{2^{-N}}{L_N} \text{ for any } N \in \mathbb{N}. \quad (35)$$

Select a continuous function  $\tilde{g}_G$  such that  $\tilde{g}_G(x) = f_G(x)$  for  $x \in U_G$ ,  $\tilde{g}_G(x) = 0$  if  $x \notin \tilde{U}_G$  and  $|\tilde{g}_G| \leq 1$ . Hence  $\tilde{g}_G \geq f_G$  on  $\mathbb{R}$ , and  $D(\tilde{g}_G, \Lambda) \supset D(f_G, \Lambda) \supset G$ .

It is also clear that  $0 \leq \tilde{g}_G - f_G \leq \chi_{\tilde{U}_G \setminus U_G} =: h_G$ , and

$$\sum_{\lambda \in \Lambda} \left( \tilde{g}_G(x + \lambda) - f_G(x + \lambda) \right) \leq \sum_{\lambda \in \Lambda} h_G(x + \lambda). \quad (36)$$

Next we prove that

$$\sum_{\lambda \in \Lambda} h_G(x + \lambda) \text{ is finite almost everywhere,} \quad (37)$$

yielding that  $C(\tilde{g}_G, \Lambda)$  equals  $C(f_G, \Lambda)$  modulo a set of measure zero.

Put  $H_{G,K,\infty} = \{x \in [-K, K] : \sum_{\lambda \in \Lambda} h_G(x + \lambda) = \infty\}$ . We will show that

$$\text{for any } K > 1 \text{ we have } \mu(H_{G,K,\infty}) = 0. \quad (38)$$

This clearly implies (37).

Observe that if  $x \in H_{G,K,\infty}$ , then there are infinitely many  $\lambda$ s such that  $x + \lambda \in \tilde{U}_G \setminus U_G$ , that is,  $x \in ((\tilde{U}_G \setminus U_G) - \lambda) \cap [-K, K]$ . Thus, by the Borel–Cantelli lemma to prove (38) it is sufficient to show that

$$\sum_{\lambda \in \Lambda} \mu\left(\left((\tilde{U}_G \setminus U_G) - \lambda\right) \cap [-K, K]\right) < \infty. \quad (39)$$

This is shown by the following estimate

$$\begin{aligned} \sum_{\lambda \in \Lambda} \mu\left(\left((\tilde{U}_G \setminus U_G) - \lambda\right) \cap [-K, K]\right) &= \sum_{\lambda \in \Lambda} \sum_{N=1}^{\infty} \mu\left(\left(\left((\tilde{U}_G \setminus U_G) \cap [N-1, N]\right) - \lambda\right) \cap [-K, K]\right) \\ &= \sum_{N=1}^{\infty} \sum_{\lambda \in \Lambda} \mu\left(\left(\left((\tilde{U}_G \setminus U_G) \cap [N-1, N]\right) \cap [\lambda - K, \lambda + K]\right)\right) \\ &= \sum_{N=1}^K \sum_{\lambda \in \Lambda} \mu\left(\left(\left((\tilde{U}_G \setminus U_G) \cap [N-1, N]\right) \cap [\lambda - K, \lambda + K]\right)\right) \\ &\quad + \sum_{N=K+1}^{\infty} \sum_{\lambda \in \Lambda} \mu\left(\left(\left((\tilde{U}_G \setminus U_G) \cap [N-1, N]\right) \cap [\lambda - K, \lambda + K]\right)\right) \end{aligned}$$

(with a finite  $S_1$ )

$$= S_1 + \sum_{N=K+1}^{\infty} \sum_{\lambda \in \Lambda, \lambda \leq 10N} \mu\left(\left(\left((\tilde{U}_G \setminus U_G) \cap [N-1, N]\right) \cap [\lambda - K, \lambda + K]\right)\right)$$

(now using (34) and (35))

$$\leq S_1 + \sum_{N=K+1}^{\infty} L_N \cdot \frac{2^{-N}}{L_N} < \infty.$$

So far we have shown that  $\tilde{g}_G$  satisfies (1) and (2). Since  $\tilde{g}_G \in C^+(\mathbb{R})$ , but not in  $C_0^+(\mathbb{R})$ . We need to adjust it a little further.

Since  $G$  is open choose an increasing sequence of compact sets  $G_K \subset G \cap [-K, K]$  such that  $\bigcup_{K=1}^{\infty} G_K = G$ .

Put  $M_0 = 0$ . Choose  $M_1 \in \mathbb{R}$  such that for any  $x \in G_1$  we have

$$\sum_{\lambda \in \Lambda, M_0+10 < \lambda < M_1} \tilde{g}_G(x + \lambda) > 1,$$

and  $\tilde{g}_G(M_1 + 5) = 0$ . This latter property can be satisfied since by assumption  $\tilde{U}_G$  does not contain a half-line.

In general, if we already have selected  $M_{K-1}$  such that  $\tilde{g}_G(M_{K-1} + 5(K-1)) = 0$  then choose  $M_K \in \mathbb{R}$  such that for any  $x \in G_K$  we have

$$\sum_{\lambda \in \Lambda, M_{K-1}+10K < \lambda < M_K} \tilde{g}_G(x + \lambda) > K, \quad (40)$$

and  $\tilde{g}_G(M_K + 5K) = 0$ .

For  $x \leq M_1 + 5$  we put  $g_G(x) = \tilde{g}_G(x)$ . For  $K > 1$  and  $x \in (M_{K-1} + 5(K-1), M_K + 5K]$  we put  $g_G(x) = \frac{1}{K} \tilde{g}_G(x)$ .

It is clear that  $g_G \in C_0^+(\mathbb{R})$ .

Since  $g_G \leq \tilde{g}_G$  we have  $C(g_G, \Lambda) \supset C(\tilde{g}_G, \Lambda)$ . If we can show that  $G \subset D(g_G, \Lambda)$  then we are done. Suppose  $x \in G$ . Then there is a  $K_x$  such that  $x \in G_{K_x}$  for any  $K \geq K_x$ . Therefore, for these  $K$  we have  $x \in [-K_x, K_x] \subset [-K, K]$  and by using (40)


$$\sum_{\lambda \in \Lambda, M_{K-1}+6K < \lambda < M_K+4K} g_G(x + \lambda) = \sum_{\lambda \in \Lambda, M_{K-1}+6K < \lambda < M_K+4K} \frac{1}{K} \tilde{g}_G(x + \lambda) > 1,$$

for any  $K \geq K_x$  and hence  $x \in D(g_G, \Lambda)$ .  $\square$

### 3 Subintervals in $C(f, \Lambda)$

**Theorem 3.1.** *There exists an asymptotically dense infinite discrete set  $\Lambda$  such that for any open set  $G \subset \mathbb{R}$  one can select an  $f_G \in C_0^+(\mathbb{R})$  such that  $D(f, \Lambda) = G$ .*

**Remark 3.2.** As Theorem 3.4 shows in the above theorem we cannot assume that  $\Lambda$  is a decreasing gap set. On the other hand, in our claim we have  $D(f, \Lambda) = G$ , that is, there is no exceptional set of measure zero where we do not know what happens. This also implies that if the interior of  $\mathbb{R} \setminus G$  is non-empty then  $C(f, \Lambda)$  contains intervals.



**Figure 2:** Definition of  $I_j$ ,  $U_j$  and related sets

*Proof.* Denote by  $\mathcal{I}_D = \{[(k-1)/2^l, k/2^l] : k, l \in \mathbb{Z}, l \geq 0\}$  the system of dyadic intervals. It is clear that one can enumerate the elements of  $\mathcal{I}_D$  in a sequence  $\{I_j\}_{j=1}^\infty$  which satisfies the following properties

$$I_j = [a_{I_j}, b_{I_j}] = \left[ \frac{k_j - 1}{2^{l_j}}, \frac{k_j}{2^{l_j}} \right] \subset [-j, j] \text{ and } \mu(I_j) = 2^{-l_j} \geq \frac{1}{j}. \quad (41)$$

We denote by  $\bar{I}_j$  the closed interval which is concentric with  $I_j$  but is of length three times the length of  $I_j$ .

We put

$$U_j = [a_j, b_j] = [2^j, 2^j + 2^{-2^j}] \text{ and } \bar{U}_j = [a_j - 2^{-2^j - j - 1}, b_j + 2^{-2^j - j - 1}] = [\bar{a}_j, \bar{b}_j].$$

See Figure 2.

We suppose that  $f_j(x) = 0$  if  $x \notin \bar{U}_j$ ,  $f_j(x) = 2^{-j}$  if  $x \in U_j$ , the function  $f_j$  is continuous on  $\mathbb{R}$  and is linear on the connected components of  $\bar{U}_j \setminus U_j$ . We define

$$\begin{aligned} \Lambda_{1,j} &= \{k \cdot 2^{-2^j - j} : k \in \mathbb{Z}\} \cap [2^j - k_j 2^{-l_j}, 2^j + 2^{-2^j} - (k_j - 1)2^{-l_j}] \\ &= \{k \cdot 2^{-2^j - j} : k \in \mathbb{Z}\} \cap [a_j - b_{I_j}, b_j - a_{I_j}] \end{aligned} \quad (42)$$

and put  $\Lambda_1 = \bigcup_{j=1}^\infty \Lambda_{1,j}$ .

Observe that if  $x \in I_j$  then

$$x + \min \Lambda_{1,j} \leq b_{I_j} + \min \Lambda_{1,j} = b_{I_j} + a_j - b_{I_j} = a_j$$

and

$$x + \max \Lambda_{1,j} \geq a_{I_j} + \max \Lambda_{1,j} = a_{I_j} + b_j - a_{I_j} = b_j,$$

hence

$$\sum_{\lambda \in \Lambda_{1,j}} f_j(x + \lambda) \geq \frac{\text{diam } U_j}{2^{-2^j-j}} 2^{-j} = \frac{2^{-2^j}}{2^{-2^j-j}} 2^{-j} = 1. \quad (43)$$

On the other hand, by (41)

$$\begin{aligned} \bar{U}_j - \Lambda_{1,j} &= [\min \bar{U}_j - \max \Lambda_{1,j}, \max \bar{U}_j - \min \Lambda_{1,j}] \\ &= [\bar{a}_j - b_j + a_{I_j}, \bar{b}_j - a_j + b_{I_j}] = [a_{I_j} - 2^{-2^j} - 2^{-2^j-j-1}, b_{I_j} + 2^{-2^j} + 2^{-2^j-j-1}] \\ &\subset \left[ a_{I_j} - \frac{1}{j}, b_{I_j} + \frac{1}{j} \right] \subset [a_{I_j} - 2^{-l_j}, b_{I_j} + 2^{-l_j}] = \bar{I}_j \end{aligned}$$

thus

$$\sum_{\lambda \in \Lambda_{1,j}} f_j(x + \lambda) = 0 \text{ if } x \in [-j, j], x \notin \bar{I}_j. \quad (44)$$

Suppose  $G \subset \mathbb{R}$  is a given open set and put  $\mathcal{J}_G = \{j : \bar{I}_j \subset G\}$ . Let  $f_G(x) = \sum_{j \in \mathcal{J}_G} f_j(x)$ . Then  $f_G$  is continuous and non-negative on  $\mathbb{R}$  and clearly  $\lim_{x \rightarrow \infty} f(x) = 0$ .

We claim that

$$\sum_{\lambda \in \Lambda_1} f_G(x + \lambda) = +\infty \quad (45)$$

exactly on  $G$ .

Indeed, if  $x \in G$  then there are infinitely many  $j$ s such that  $x \in I_j \subset \bar{I}_j \subset G$ . This means that (43) holds for infinitely many  $j \in \mathcal{J}_G$  and hence (45) is true when  $x \in G$ .

Next we need to verify that (45) does not hold for  $x \notin G$ . Suppose that  $j_0 \geq 10$ ,  $j_0 \in \mathcal{J}_G$ ,  $x \notin G$  and  $x \in [-j_0, j_0]$ . Then  $x \notin \bar{I}_{j_0}$  and by (44) we have

$$\sum_{\lambda \in \Lambda_{1,j_0}} f_{j_0}(x + \lambda) = 0. \quad (46)$$

Next assume that  $j < j_0$ . Then by using (41) and (42)

$$\begin{aligned} \max\{x + \lambda : \lambda \in \Lambda_{1,j}\} &\leq j_0 + 2^j + 2^{-2^j} - (k_j - 1)2^{-l_j} \leq j_0 + 2^j + 2^{-2^j} + j \\ &< 2j_0 + 2^{j_0-1} + 1 < 2^{j_0} - 1 < 2^{j_0} - 2^{-2^{j_0}-j_0-1} = \bar{a}_{j_0}. \end{aligned}$$

Hence,

$$\sum_{\lambda \in \Lambda_{1,j}} f_{j_0}(x + \lambda) = 0. \quad (47)$$

If  $j_0 < j$  then

$$\min\{x + \lambda : \lambda \in \Lambda_{1,j}\} \geq -j_0 + 2^j - j > 2^{j-1} - 2j - 1 + 2^{j-1} + 1 > 2^{j_0} + 1 > \bar{b}_{j_0},$$



and hence in this case we also have (47).

Therefore, from (46) and (47) it follows that

$$\sum_{\lambda \in \Lambda_1} f_{j_0}(x + \lambda) = 0 \text{ for } j_0 \in \mathcal{J}_G, j_0 \geq 10, |x| \leq j_0. \quad (48)$$

This implies

$$\sum_{\lambda \in \Lambda_1} f_G(x + \lambda) \leq \sum_{\substack{\lambda \in \Lambda_{1,j} \\ j \leq \max\{10, |x|\}}} f_j(x + \lambda) < +\infty.$$

Since  $\Lambda_1$  is not asymptotically dense we need to choose an asymptotically dense  $\Lambda_2$  such that

$$\sum_{\lambda \in \Lambda_2} \sum_{j=1}^{\infty} f_j(x + \lambda) < +\infty \text{ holds for any } x \in \mathbb{R}. \quad (49)$$

Then for any open  $G \subset \mathbb{R}$

$$\sum_{\lambda \in \Lambda_2} f_G(x + \lambda) \leq \sum_{\lambda \in \Lambda_2} \sum_{j=1}^{\infty} f_j(x + \lambda) < +\infty$$

holds and if we let  $\Lambda = \Lambda_1 \cup \Lambda_2$  then  $\Lambda$  is asymptotically dense and  $D(f_G, \Lambda) = G$ .

To complete the proof of this theorem we need to verify (49) for a suitable  $\Lambda_2$ . For  $j \geq 10$  put

$$\Lambda_{2,j} = \{k \cdot 2^{-j} : k \in \mathbb{Z}\} \cap (2^{j-1} + 2(j-1), 2^j + 2j], \text{ and } \Lambda_2 = \bigcup_{j=10}^{\infty} \Lambda_{2,j}.$$

Suppose  $x \in [-j_0, j_0]$  and  $j_0 \geq 10$ . Then for  $j \geq j_0$  from  $x + \lambda \in \overline{U}_j$  it follows that  $2^j - 1 < x + \lambda \leq j + \lambda$ , and hence

$$\lambda > 2^j - j - 1 > 2^{j-1} + 2(j-1).$$

Similarly,  $x + \lambda \in \overline{U}_j$  implies  $2^j + 1 > x + \lambda \geq -j + \lambda$ , and hence

$$\lambda < 2^j + j + 1 < 2^j + 2j.$$

Thus from  $x + \lambda \in \overline{U}_j$  it follows that  $\lambda \in \Lambda_{2,j}$ . Since the length of  $\overline{U}_j$  is less than  $2 \cdot 2^{-2^j} < 2^{-j}$  there is at most one  $\lambda \in \Lambda_{2,j}$  for which  $f_j(x + \lambda) \neq 0$  and for this  $\lambda$  we have  $f_j(x + \lambda) = 2^{-j}$ .

Put  $M_x = \max\{10, |x|\}$ . Then

$$\sum_{\lambda \in \Lambda_2} \sum_{j=1}^{\infty} f_j(x + \lambda) = \sum_{\lambda \in \Lambda_2} \sum_{j=1}^{M_x} f_j(x + \lambda) + \sum_{j=M_x+1}^{\infty} \sum_{\lambda \in \Lambda_2} f_j(x + \lambda)$$

$$\leq \sum_{\lambda \in \Lambda_2} \sum_{j=1}^{M_x} f_j(x + \lambda) + \sum_{j=M_x+1}^{\infty} 2^{-j} < +\infty.$$

□

In Theorem 2.1 we verified that for decreasing gap asymptotically dense sets  $D(f, \Lambda)$  can contain an open set, while  $C(f, \Lambda)$  equals the complement of this open set only almost everywhere.

The next example shows that one can define decreasing gap asymptotically dense  $\Lambda$ s for which one can find nonnegative continuous  $f$ s such that both  $C(f, \Lambda)$  and  $D(f, \Lambda)$  have interior points.

**Theorem 3.3.** *There exists a decreasing gap asymptotically dense  $\Lambda$  and an  $f \in C_0^+(\mathbb{R})$  such that  $I_1 = [0, 1] \subset D(f, \Lambda)$  and  $I_2 = [4, 5] \subset C(f, \Lambda)$ .*

*Proof.* Put  $f(x) = 2^{-2^{j+1}}$  if  $x \in [10j, 10j + 1]$  for a  $j \in \mathbb{N}$ . Set  $f(x) = 0$  if  $x \in \{10j - 1/4, 10j + 5/4\}$  for a  $j \in \mathbb{N}$ , and also put  $f(x) = 0$  for  $x \leq 0$ . We suppose that  $f$  is linear on the intervals where we have not defined it so far. Put  $\Lambda_{1,j} = \{k \cdot 2^{-2^j} : k \in \mathbb{Z}\} \cap [10j - 10, 10j - 2)$  and  $\Lambda_{2,j} = \{k \cdot 2^{-2^{j+1}} : k \in \mathbb{Z}\} \cap [10j - 2, 10j)$ . Let  $\Lambda = \bigcup_{j=1}^{\infty} (\Lambda_{1,j} \cup \Lambda_{2,j})$ . Observe that  $\Lambda$  is a decreasing gap asymptotically dense set.

One can see that for  $x \in I_1$  we have

$$\sum_{\lambda \in \Lambda} f(x + \lambda) \geq \sum_{j=1}^{\infty} 2^{2^{j+1}} \cdot 2^{-2^{j+1}} = +\infty$$

and for  $x \in I_2$

$$\sum_{\lambda \in \Lambda} f(x + \lambda) \leq \sum_{j=1}^{\infty} 2 \cdot 2^{2^j} \cdot 2^{-2^{j+1}} < +\infty.$$

It is also clear from the construction that  $\lim_{x \rightarrow \infty} f(x) = 0$ . □

Observe that in the above construction  $I_1 \subset D(f, \Lambda)$  was to the left of  $I_2 \subset C(f, \Lambda)$ . The next theorem shows that for decreasing gap asymptotically dense  $\Lambda$ s and continuous functions this situation cannot be improved. If  $x$  is an interior point of  $C(f, \Lambda)$  then the half-line  $[x, \infty)$  intersects  $D(f, \Lambda)$  in a set of measure zero. As Theorem 3.1 shows if we do not assume that  $\Lambda$  is of decreasing gap then it is possible that  $D(f, \Lambda)$  has a part of positive measure, even to the right of the interior points of  $C(f, \Lambda)$ .

**Theorem 3.4.** *Let  $\Lambda$  be a decreasing gap and asymptotically dense set, and let  $f : \mathbb{R} \rightarrow [0, +\infty)$  be continuous. Then if  $x$  is an interior point of  $C(f, \Lambda)$  then*

$$\mu\left([x, +\infty) \cap D(f, \Lambda)\right) = 0. \quad (50)$$

*Proof.* Proceeding towards a contradiction assume the existence of a non-degenerate closed interval  $I \subset C(f, \Lambda)$ . Suppose that there is a bounded subset  $D_1(f, \Lambda) \subset D(f, \Lambda)$  with positive measure to the right of  $I$ . Choose an interval  $J = [a_J, b_J]$  to the right of  $I$  such that

$$\mu(J) = \mu(I)/10, \text{ and } \mu(J \cap D(f, \Lambda)) = \alpha > 0. \quad (51)$$

We put  $D_1(f, \Lambda) = J \cap D(f, \Lambda)$ . We suppose that  $\Lambda = \{\lambda_1, \lambda_2, \dots\}$  is indexed in an increasing order. Select  $N$  such that

$$\lambda_n - \lambda_{n-1} < \frac{\mu(I)}{100} \text{ for } n \geq N. \quad (52)$$

We clearly have that  $\sum_{i=N}^{\infty} f(x + \lambda_i)$  diverges on  $D_1(f, \Lambda)$ . Moreover, if  $n \in \mathbb{N}$ , which is to be fixed later, for large enough  $M$  we have  $\sum_{i=N}^M f(x + \lambda_i) > n$  in a set  $D_2(f, \Lambda) \subset D_1(f, \Lambda)$  of measure larger than  $\frac{\alpha}{2}$ . Hence we have

$$\int_{D_2(f, \Lambda)} \sum_{i=N}^M f(x + \lambda_i) dx \geq \frac{n\alpha}{2}. \quad (53)$$

Assume that  $i \in \{N, N+1, \dots, M\}$ . We choose  $\gamma(i)$  such that

$$a_J + \lambda_i - \lambda_{\gamma(i)} \in I, \text{ but } a_J + \lambda_i - \lambda_{\gamma(i)+1} \notin I. \quad (54)$$

Since  $a_J$  is to the right of  $I$  it is clear that  $\lambda_{\gamma(i)} > \lambda_i$ , therefore  $\gamma(i) > i \geq N$  and hence (52) implies that  $\gamma(i)$  is well-defined, that is (54) can be satisfied.

It is also clear that there exists  $\widetilde{M}$  such that  $\gamma(i) \leq \widetilde{M}$  holds for  $i \in \{N, N+1, \dots, M\}$ .

By (51), (52), and (54) we have

$$J + \lambda_i - \lambda_{\gamma(i)} \subset I \text{ and hence } D_2(f, \Lambda) + \lambda_i - \lambda_{\gamma(i)} \subset I. \quad (55)$$

Next we verify that

$$\text{if } i' \neq i \text{ then } \gamma(i') \neq \gamma(i). \quad (56)$$

Indeed, we can suppose that  $i' < i$ , and proceeding towards a contradiction we also suppose that  $\gamma(i') = \gamma(i)$ . We know that  $a_J + \lambda_i - \lambda_{\gamma(i)} \in I$ , moreover  $a_J + \lambda_{i'} - \lambda_{\gamma(i')} \in I$  holds as well. Since  $\gamma(i) = \gamma(i')$  we have

$$a_J + \lambda_{i'} - \lambda_{\gamma(i')} = a_J + \lambda_i - \lambda_{\gamma(i)} - \lambda_i + \lambda_{i'} \in I.$$

Using the first half of (54) and  $\lambda_{i'} \leq \lambda_{i-1} < \lambda_i$  we also obtain

$$a_J + \lambda_i - \lambda_{\gamma(i)} - \lambda_i + \lambda_{i'} \leq a_J + \lambda_i - \lambda_{\gamma(i)} - \lambda_i + \lambda_{i-1} \in I.$$

Since  $\Lambda$  is of decreasing gap and  $\gamma(i) > i$  we have  $\lambda_{\gamma(i)+1} - \lambda_{\gamma(i)} < \lambda_i - \lambda_{i-1}$ , and hence

$$a_J + \lambda_i - \lambda_{\gamma(i)} - \lambda_i + \lambda_{i-1} < a_J + \lambda_i - \lambda_{\gamma(i)} - \lambda_{\gamma(i)+1} + \lambda_{\gamma(i)} \in I,$$

which contradicts (54).

By using (55) and (56) we infer

$$\begin{aligned} \int_{D_2(f,\Lambda)} \sum_{i=N}^M f(x + \lambda_i) dx &= \sum_{i=N}^M \int_{D_2(f,\Lambda)} f(x + \lambda_i - \lambda_{\gamma(i)} + \lambda_{\gamma(i)}) dx \quad (57) \\ &= \sum_{i=N}^M \int_{D_2(f,\Lambda) + \lambda_i - \lambda_{\gamma(i)}} f(t + \lambda_{\gamma(i)}) dt \leq \int_I \sum_{j=N}^{\tilde{M}} f(t + \lambda_j) dt. \end{aligned}$$

Thus by (53) we obtain

$$\int_I \sum_{i=N}^{\tilde{M}} f(x + \lambda_i) dx \geq \frac{n\alpha}{2},$$

as the left-handside by (57) gives an upper bound for the integral in (53). However,  $\sum_{i=N}^{\tilde{M}} f(x + \lambda_i)$  is continuous, which yields that this integrand is at least  $\frac{n\alpha}{4\mu(I)}$  in a non-degenerate closed subinterval  $I_1 \subset I$ . Thus we have  $s(x) = \sum_{\lambda \in \Lambda} f(x + \lambda) > \frac{n\alpha}{4\mu(I)}$  in  $I_1$ . Hence, if we choose  $n$  to be large enough, we find that  $s(x) > 1$  in  $I_1$ .

Now by applying the very same argument to  $I_1$  instead of  $I$ , we might obtain that  $s(x) > \frac{n_1\alpha}{4\mu(I_1)}$  in a non-degenerate closed subinterval  $I_2 \subset I_1$ . Thus if we choose  $n_1$  to be large enough, we find that  $s(x) > 2$  in  $I_2$ . Proceeding recursively we obtain a nested sequence of closed intervals  $I_1, I_2, \dots$  such that  $s(x) > k$  for  $x \in I_k$ . As this system of intervals has a nonempty intersection, we find that there is a point in  $I$  with  $s(x) = \infty$ , a contradiction.  $\square$

## 4 Acknowledgements

During the Fall semester of 2018, when this paper was prepared all three authors visited the Institut Mittag-Leffler in Djursholm and participated in the semester Fractal Geometry and Dynamics. We thank the hospitality and financial support of the Institut Mittag-Leffler. Z. Buczolich also thanks the Rényi Institute where he was a visiting researcher for the academic year 2017-18.

## References

- [1] Z. Buczolich, J-P. Kahane and R. D. Mauldin, On series of translates of positive functions, *Acta Math. Hungar.*, **93**(3) (2001), 171-188.
- [2] Z. Buczolich, J-P. Kahane, and R. D. Mauldin, Sur les séries de translatées de fonctions positives. *C. R. Acad. Sci. Paris Sér. I Math.*, **329**(4):261–264, 1999.
- [3] Z. Buczolich and R. D. Mauldin, On the convergence of  $\sum_{n=1}^{\infty} f(nx)$  for measurable functions. *Mathematika*, **46**(2):337–341, 1999.
- [4] Z. Buczolich and R. D. Mauldin, On series of translates of positive functions II., *Indag. Mathem., N. S.*, **12** (3), (2001), 317-327.
- [5] J.A. Haight, A linear set of infinite measure with no two points having integral ratio, *Mathematika* **17**(1970), 133-138.
- [6] J.A. Haight, A set of infinite measure whose ratio set does not contain a given sequence, *Mathematika* **22**(1975), 195-201.
- [7] C. G. Lekkerkerker, Lattice points in unbounded point sets, I. *Indag. Math.*, **20** (1958) 197-205.
- [8] H. v. Weizsäcker, *Zum Konvergenzverhalten der Reihe  $\sum_{n=1}^{\infty} f(nt)$  für  $\lambda$ -messbare Funktionen  $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$* , Diplomarbeit, Universität München, 1970.